

FORBIDDEN GAP ARGUMENT FOR PHASE TRANSITIONS PROVED BY MEANS OF CHESSBOARD ESTIMATES

MAREK BISKUP¹ AND ROMAN KOTECKÝ²

¹*Department of Mathematics, UCLA, Los Angeles, California, USA*

²*Center for Theoretical Study, Charles University, Prague, Czech Republic*

Abstract: Existence of first-order phase transitions is often proved with the aid of reflection positivity and chessboard estimates. The standard approach relies on estimates of correlations in torus measures which yield the existence of a transition point where the free energy has a discontinuous derivative with respect to a suitably chosen variable. In addition, at the transition point, two distinct translation-invariant Gibbs states are extracted from torus measures in which the one-sided derivatives of the free energy are realized as expectations of a local observable X . Here we show that (most of) the gap between these extreme expected values is forbidden: There are no shift-ergodic Gibbs states for which the expectation of X lies deep inside the gap. We point out several recent results based on chessboard estimates where our main theorems provide important additional information concerning the structure of the set of possible thermodynamic equilibria.

1. INTRODUCTION

One of the basic tasks of mathematical statistical mechanics is to find a rigorous approach to various first-order phase transitions in lattice spin systems. Here, two methods of proof are generally available: Pirogov-Sinai theory and chessboard estimates. The former, developed in [19, 20], possesses an indisputable advantage of robustness with respect to (general) perturbations, but its drawback is the (nearly strict) restriction to finite sets of possible spin values. The latter method, which goes back to [12, 13], is limited, for the most part, to systems with nearest-neighbor attractive interactions but it poses almost no limitations on the individual spin space. In particular, the method of chessboard estimates is still the only tool we have for a general control of asymmetric phase transitions in continuum spin systems.

A closer look at the results proved using the aforementioned methods reveals one important additional difference. Namely, while both techniques ultimately produce a proof of phase coexistence, Pirogov-Sinai theory offers significantly better control of the number of possible Gibbs states. In fact, one can prove the so called *completeness of phase diagram* [22] which essentially asserts that the states constructed by the theory exhaust the set of all shift-ergodic Gibbs states. In a more technical language, there is a one-to-one correspondence between the shift-ergodic Gibbs states and the “stable phases” defined in terms of minimal “metastable free energy”. No statement of this kind has been available in the approaches based solely on chessboard estimates. This

© 2005 by M. Biskup and R. Kotecký. Reproduction, by any means, of the entire article for non-commercial purposes is permitted without charge.

makes many of the conclusions based on this technique—see [2, 4, 8, 11, 21] for a modest sample of recent references—seem to be somewhat “incomplete.”

To make the distinction more explicit, let us consider the example of temperature-driven first-order phase transition in the q -state Potts model with $q \gg 1$. In dimensions $d \geq 2$, there exists a transition temperature, T_t , at which there are q ordered states—related by an underlying symmetry of the model—that are low on both entropy and energy, and one disordered state which is abundant in both quantities. The transition is accompanied by a massive jump in the energy density (as a function of temperature). Here the proof based on chessboard estimates [16] seems to produce “only” the existence of a temperature where the aforementioned $q + 1$ states coexist, but it does not rule out the existence of other states; particularly, those with energies “inside” the jump. On the other hand, Pirogov-Sinai approaches [15, 17] permit us to conclude that *no other* than the above $q + 1$ shift-ergodic Gibbs states can exist at T_t and, in particular, there is a *forbidden gap* of energy densities where no shift ergodic Gibbs states are allowed to enter.

The purpose of this note is to show that, after all, chessboard estimates can also be supplemented with a corresponding forbidden-gap argument. As a consequence, the theorems of, e.g., [2, 4, 8, 11, 16, 21] imply not only the *existence* of particular states at the corresponding transition temperature, but also the *absence* of states that differ significantly from those proved to exist. Our arguments are fairly general; indeed, apart from the necessary condition of reflection positivity we require only translation invariance and absolute summability of interactions. The main idea of the proof is that all Gibbs states (at the same temperature) have the same large-deviation properties on the scale that is exponential in volume. This permits us to compare any translation-invariant Gibbs state with a corresponding measure on torus, where chessboard estimates can be used to rule out most of undesirable scenarios.

The rest of this paper is organized as follows: In Sect. 2.1 and 2.2 we define the class of models to which our techniques apply and review various elementary facts about reflection positivity and chessboard estimates. The statements of our main theorems (Theorem 2.5 and Corollary 2.6) come in Sect 2.3. The proofs constitute the bulk of Sect. 3; applications to recent results established by means of chessboard estimates are discussed in Sect. 4. The Appendix (Sect. 5) contains the proof of Theorem 4.4 which is a refined version of Theorem 3 of [11]. This result is needed for one of our applications in Sect. 4.

2. MAIN RESULT

In order to formulate our principal claims we will first recall the standard setup for proofs of first-order phase transitions by chessboard estimates and introduce the necessary notations. The actual theorems are stated in Sect. 2.3.

2.1 Models of interest.

We will work with the standard class of spin systems on \mathbb{Z}^d and so we will keep our discussion of general concepts at the minimum possible. We refer the reader to Georgii’s monograph [14] for a more comprehensive treatment and relevant references.

Our spins, s_x , will take values in a compact separable metric space Ω_0 . We equip Ω_0 with the σ -algebra \mathcal{F}_0 of its Borel subsets and consider an *a priori* probability measure ν_0 on $(\Omega_0, \mathcal{F}_0)$. Spin configurations on \mathbb{Z}^d are the collections $(s_x)_{x \in \mathbb{Z}^d}$. We will use $\Omega = \Omega_0^{\mathbb{Z}^d}$ to denote the set of all spin configurations on \mathbb{Z}^d and \mathcal{F} to denote the σ -algebra of Borel subsets of Ω defined using the product topology. If $\Lambda \subset \mathbb{Z}^d$, we define \mathcal{F}_Λ to be the sub- σ -algebra of events depending only on $(s_x)_{x \in \Lambda}$. For each $x \in \mathbb{Z}^d$, the map $\tau_x: \Omega \rightarrow \Omega$ is the “translation by x ” defined by $(\tau_x s)_y = s_{x+y}$. It is easy to check that τ_x is a continuous and hence measurable for all $x \in \mathbb{Z}^d$. We will write $\Lambda \Subset \mathbb{Z}^d$ to indicate that Λ is a finite subset of \mathbb{Z}^d .

To define Gibbs measures, we will consider a family of Hamiltonians $(H_\Lambda)_{\Lambda \Subset \mathbb{Z}^d}$. These will be defined in terms of interaction potentials $(\Phi_A)_{A \Subset \mathbb{Z}^d}$. Namely, for each $A \Subset \mathbb{Z}^d$, let $\Phi_A: \Omega \rightarrow \mathbb{R}$ be a function with the following properties:

- (1) The function Φ_A is \mathcal{F}_A -measurable for each $A \Subset \mathbb{Z}^d$.
- (2) The interaction (Φ_A) is translation invariant, i.e., $\Phi_{A+x} = \Phi_A \circ \tau_x$ for all $x \in \mathbb{Z}^d$ and all $A \Subset \mathbb{Z}^d$.
- (3) The interaction (Φ_A) is absolutely summable in the sense that

$$\|\Phi\| = \sum_{\substack{A \Subset \mathbb{Z}^d \\ 0 \in A}} \|\Phi_A\|_\infty < \infty. \quad (2.1)$$

The Hamiltonian on a set $\Lambda \Subset \mathbb{Z}^d$ is a function $H_\Lambda: \Omega \rightarrow \mathbb{R}$ defined by

$$H_\Lambda = \sum_{\substack{A \Subset \mathbb{Z}^d \\ A \cap \Lambda \neq \emptyset}} \Phi_A. \quad (2.2)$$

For each $\beta \geq 0$, let \mathfrak{G}_β be the set of Gibbs measures for the Hamiltonian (2.2). Specifically, $\mu \in \mathfrak{G}_\beta$ if and only if the conditional probability $\mu(\cdot | \mathcal{F}_{\Lambda^c})$ —which exists since Ω is a Polish space—satisfies, for all $\Lambda \Subset \mathbb{Z}^d$ and μ -almost all s , the (conditional) DLR equation

$$\mu(ds_\Lambda | \mathcal{F}_{\Lambda^c})(s) = \frac{e^{-\beta H_\Lambda(s)}}{Z_\Lambda} \prod_{x \in \Lambda} \nu_0(ds_x). \quad (2.3)$$

Here $Z_\Lambda = Z_\Lambda(\beta, s_{\Lambda^c})$ is a normalization constant which is independent of $s_\Lambda = (s_x)_{x \in \Lambda}$.

Remark 2.1 The results of the present paper can be generalized even to the situations with unbounded spins and interactions; see Theorem 4.5. However, the general theory of Gibbs measures with unbounded spins features some unpleasant technicalities that would obscure the presentation. We prefer to avoid them and to formulate the bulk of the paper for systems with compact spins. Our restriction to translation-invariant interaction in (2) above is mostly for convenience of exposition. Actually, the proofs in Sect. 3 can readily be modified to include periodic interactions as well.

2.2 Chessboard estimates.

As alluded to before, chessboard estimates are one of the principal tool for proving phase coexistence. In order to make this tool available, we have to place our spin system on torus. Let \mathbb{T}_L be the torus of $L \times \cdots \times L$ sites and let $H_L: \Omega_0^{\mathbb{T}_L} \rightarrow \mathbb{R}$ be the function defined as follows.

Given a configuration $s = (s_x)_{x \in \mathbb{T}_L}$, we extend s periodically to a configuration \bar{s} on all of \mathbb{Z}^d . Using $H_{\mathbb{T}_L}$ to denote the Hamiltonian associated with the embedding of \mathbb{T}_L into \mathbb{Z}^d , we define $H_L(s) = H_{\mathbb{T}_L}(\bar{s})$. The *torus measure* $\mathbb{P}_{L,\beta}$ then simply is

$$\mathbb{P}_{L,\beta}(ds) = \frac{e^{-\beta H_L(s)}}{Z_L} \prod_{x \in \mathbb{T}_L} \nu_0(ds_x). \quad (2.4)$$

Here $Z_L = Z_L(\beta)$ is the torus partition function.

Chessboard estimates will be implied by the condition of *reflection positivity*. While this condition can already be defined in terms of interactions $(\Phi_\Lambda)_{\Lambda \in \mathbb{Z}^d}$, it is often easier to check it directly on torus. Let us consider a torus \mathbb{T}_L with even L and let us split it into two symmetric halves, \mathbb{T}_L^+ and \mathbb{T}_L^- , sharing a “plane of sites” on their boundary. We will refer to the set $P = \mathbb{T}_L^+ \cap \mathbb{T}_L^-$ as a *plane of reflection*. Let \mathcal{F}_P^+ and \mathcal{F}_P^- denote the σ -algebras of events depending only on configurations in \mathbb{T}_L^+ and \mathbb{T}_L^- , respectively.

We assume that the naturally-defined (spatial) reflection $\vartheta_P : \mathbb{T}_L^+ \leftrightarrow \mathbb{T}_L^-$ gives rise to a map $\theta_P : \Omega_0^{\mathbb{T}_L} \rightarrow \Omega_0^{\mathbb{T}_L}$ which obeys the following constraints:

- (1) θ_P is an *involution*, $\theta_P \circ \theta_P = \text{id}$.
- (2) θ_P is a *reflection* in the sense that if $\mathcal{A} \in \mathcal{F}_P^+$ depends only on configurations in $\Lambda \subset \mathbb{T}_L^+$, then $\theta_P(\mathcal{A}) \in \mathcal{F}_P^-$ depends only on configurations in $\vartheta_P(\Lambda)$.

In many cases of interest, θ_P is simply the mapping that is directly induced by the spacial reflection ϑ_P , i.e., $\theta_P = \vartheta_P^*$, where $(\vartheta_P^*(s))_x = s_{\vartheta_P(x)}$; our definition permits us to combine the spatial reflection with an involution of the single-spin space.

Reflection positivity is now defined as follows:

Definition 2.2 Let \mathbb{P} be a probability measure on $\Omega_0^{\mathbb{T}_L}$ and let \mathbb{E} be the corresponding expectation. We say that \mathbb{P} is *reflection positive*, if for any plane of reflection P and any two bounded \mathcal{F}_P^+ -measurable random variables X and Y the following inequalities hold:

$$\mathbb{E}(X\theta_P(Y)) = \mathbb{E}(Y\theta_P(X)) \quad (2.5)$$

and

$$\mathbb{E}(X\theta_P(X)) \geq 0. \quad (2.6)$$

Here, $\theta_P(X)$ denotes the \mathcal{F}_L^- -measurable random variable $X \circ \theta_P$.

Remark 2.3 Here are some standard examples of summable two-body interactions that are reflection positive. Consider spin systems with vector-valued spins s_x and interaction potentials

$$\Phi_{\{x,y\}} = J_{x,y} (s_x, s_y), \quad x \neq y, \quad (2.7)$$

where $J_{x,y}$ are coupling constants and (\cdot, \cdot) denotes a positive-semidefinite inner product on Ω . Then the corresponding torus Gibbs measure with $\beta \geq 0$ is reflection positive for the following choices of $J_{x,y}$'s:

- (1) *Nearest and next-nearest neighbor couplings*: $J_{x,y} = \lambda$ if x and y are nearest neighbors, $J_{x,y} = \kappa$ with $\lambda \geq 2(d-1)|\kappa|$ if x and y are next-nearest neighbors and $J_{x,y} = 0$ in the remaining cases.

(2) *Yukawa-type potentials*:

$$J_{x,y} = e^{-\mu|x-y|_1}, \quad (2.8)$$

where $\mu > 0$ and $|x - y|_1$ is the ℓ^1 -distance between x and y .

(3) *Power-law decaying interactions*:

$$J_{x,y} = \frac{1}{|x - y|_1^s}, \quad (2.9)$$

with $s > 0$.

The proofs of these are based on the general theory developed in [12, 13]; relevant calculations can also be found in [1, Sect. 4.2].

Of course, any linear combination of the above—as well as other reflection-positive interactions—with positive coefficients is still reflection positive.

Now, we are finally getting to the setup underlying chessboard estimates. Suppose that L is an integer multiple of an (integer) number B . (To rule out various technical complications with the following theorem, we will actually always assume that L/B is a power of 2.) Let $\Lambda_B \subset \mathbb{T}_L$ be the box of $(B + 1) \times \cdots \times (B + 1)$ sites with the “lower-left” corner at the origin—we will call such box a *B-block*. We can tile \mathbb{T}_L by translates of Λ_B by B -multiples of vectors from the *factor torus*, $\tilde{\mathbb{T}} = \mathbb{T}_{L/B}$. Note that the neighboring translates of Λ_B will have a side in common. Let \mathcal{A} be an event depending only on configurations in Λ_B ; we will call such \mathcal{A} a *B-block event*. For each $\mathbf{t} \in \tilde{\mathbb{T}}$, we define the event $\theta_{\mathbf{t}}(\mathcal{A})$ as follows:

- (1) If \mathbf{t} has all components even, then $\theta_{\mathbf{t}}(\mathcal{A})$ is simply the translation of \mathcal{A} by vector $B\mathbf{t}$, i.e., $\theta_{\mathbf{t}}(\mathcal{A}) = \tau_{B\mathbf{t}}^{-1}(\mathcal{A}) = \{s \in \Omega_0^{\mathbb{T}_L} : \tau_{B\mathbf{t}}(s) \in \mathcal{A}\}$.
- (2) For the remaining $\mathbf{t} \in \tilde{\mathbb{T}}$, we first reflect \mathcal{A} through the “midplane” of Λ_B in all directions whose component of \mathbf{t} is odd, and then translate the result by $B\mathbf{t}$ as before.

Thus, $\theta_{\mathbf{t}}(\mathcal{A})$ will always depend only on configurations in the B -block $\Lambda_B + B\mathbf{t}$.

The desired consequence of reflection positivity is now stated as follows.

Theorem 2.4 (Chessboard estimate) *Let \mathbb{P} be a measure on $\Omega_0^{\mathbb{T}_L}$ which is reflection-positive with respect to θ_P . Then for any B -block events $\mathcal{A}_1, \dots, \mathcal{A}_m$ and any distinct sites $\mathbf{t}_1, \dots, \mathbf{t}_m \in \tilde{\mathbb{T}}$,*

$$\mathbb{P}\left(\bigcap_{j=1}^m \theta_{\mathbf{t}_j}(\mathcal{A}_j)\right) \leq \prod_{j=1}^m \mathbb{P}\left(\bigcap_{\mathbf{t} \in \tilde{\mathbb{T}}} \theta_{\mathbf{t}}(\mathcal{A}_j)\right)^{1/|\tilde{\mathbb{T}}|}. \quad (2.10)$$

Proof. See [12, 13]. □

The moral of this result—whose proof is nothing more than an enhanced version of the Cauchy-Schwarz inequality applied to the inner product $X, Y \mapsto \mathbb{E}(X\theta_P(Y))$ —is that the probability of any number of events factorizes, as a bound, into the product of probabilities. This is particularly useful for contour estimates; of course, provided that the word contour refers to a collection of boxes on each of which some “bad” event occurs. Indeed, by (2.10) the probability of a contour will automatically be suppressed exponentially in the number of constituting “bad” boxes.

2.3 Main theorems.

For any B -block event \mathcal{A} , we introduce the quantity

$$\mathfrak{p}_\beta(\mathcal{A}) = \lim_{L \rightarrow \infty} \left(\mathbb{P}_{L,\beta} \left(\bigcap_{\mathbf{t} \in \tilde{\mathbb{T}}} \theta_{\mathbf{t}}(\mathcal{A}) \right) \right)^{1/|\tilde{\mathbb{T}}|}, \quad (2.11)$$

with the limit taken over multiples of B . The limit exists by standard subadditivity arguments. While the definition would suggest that $\mathfrak{p}_\beta(\mathcal{A})$ is a large-deviation rate, chessboard estimates (2.10) show that $\mathfrak{p}_\beta(\mathcal{A})$ can also be thought of as the “probability of \mathcal{A} regardless of the status of all other B -blocks.” This interpretation is supported by the fact that $\mathcal{A} \mapsto \mathfrak{p}_\beta(\mathcal{A})$ is an outer measure on \mathcal{F}_{Λ_B} with $\mathfrak{p}_\beta(\Omega) = 1$, cf. Lemma 6.3 of [4].

Furthermore, recalling that Λ_{N-1} is the block of $N \times \dots \times N$ sites with the “lower-left” corner at the lattice origin, let

$$R_N(\mathcal{A}) = \frac{1}{|\Lambda_{N-1}|} \sum_{x \in \Lambda_{N-1}} \mathbf{1}_{\mathcal{A} \circ \tau_{Bx}} \quad (2.12)$$

be the fraction of B -blocks (in Λ_{N-1}) in which \mathcal{A} occurs. Whenever $\mu \in \mathfrak{G}_\beta$ is a Gibbs state for the Hamiltonian (2.2) at inverse temperature β that is invariant with respect to the shifts $(\tau_{Bx})_{x \in \mathbb{Z}^d}$, the limit

$$\rho_\mu(\mathcal{A}) = \lim_{N \rightarrow \infty} R_N(\mathcal{A}) \quad (2.13)$$

exists μ -almost surely. In the following, we will use $\rho_\mu(\mathcal{A})$ mostly for measures that are actually ergodic with respect to the shifts by multiples of B . In such cases the limit is self-averaging, $\rho_\mu(\mathcal{A}) = \mu(\mathcal{A})$ almost surely. Notwithstanding, we will stick to the notation $\rho_\mu(\mathcal{A})$ to indicate that claims are being made about almost-sure properties of configurations and not just expectations. To keep our statements concise, we will refer to measures which are invariant and ergodic with respect to the translations $(\tau_{Bx})_{x \in \mathbb{Z}^d}$ as *B -shift ergodic*.

Our principal result is now as follows:

Theorem 2.5 *Consider a spin system as described above and suppose that the torus measure is reflection positive for all $\beta \geq 0$ and all even $L \geq 2$. Let $\mathcal{G}_1, \dots, \mathcal{G}_r$ be a finite number of “good” B -block events and let $\mathcal{B} = (\mathcal{G}_1 \cup \dots \cup \mathcal{G}_r)^c$ be the corresponding “bad” B -block event. Suppose that the good block events are mutually exclusive and non-compatible (different types of goodness cannot occur in neighboring translates of Λ_B):*

- (1) $\mathcal{G}_i \cap \mathcal{G}_j = \emptyset$ for all $i \neq j$.
- (2) If $\mathbf{t}_1, \mathbf{t}_2 \in \tilde{\mathbb{T}}$ are nearest neighbors, then

$$\theta_{\mathbf{t}_1}(\mathcal{G}_i) \cap \theta_{\mathbf{t}_2}(\mathcal{G}_j) = \emptyset, \quad 1 \leq i < j \leq r. \quad (2.14)$$

Then for every $\epsilon > 0$, there exists $\delta > 0$ such that for any $\beta \geq 0$ with $\mathfrak{p}_\beta(\mathcal{B}) < \delta$ we have

$$\rho_\mu(\mathcal{B}) \in [0, \epsilon] \quad (2.15)$$

and

$$\rho_\mu(\mathcal{G}_i) \in [0, \epsilon] \cup [1 - \epsilon, 1] \quad (2.16)$$

for every B -shift ergodic Gibbs state $\mu \in \mathfrak{G}_\beta$ and all $i = 1, \dots, r$.

We remark that the conclusion of Theorem 2.5 holds even when the requirement of compact spin-spin space and norm-bounded interactions are relaxed to the condition of *finite average energy*. We state the corresponding generalization in Theorem 4.5. Theorem 2.5 directly implies the standard conclusion of chessboard estimates (cf. [9, Propositions 3.1-3.3] or [16, Theorem 4]):

Corollary 2.6 *Let $\beta_1 < \beta_2$ be two inverse temperatures and let \mathcal{G}_1 and \mathcal{G}_2 be two mutually exclusive and non-compatible good B -block events (conditions (1) and (2) in Theorem 2.5). Then, for every $\epsilon > 0$ there exists a constant $\delta > 0$ such that the conditions*

- (1) $\mathfrak{p}_\beta(\mathcal{B}) < \delta$ for all $\beta \in [\beta_1, \beta_2]$ and
- (2) $\mathfrak{p}_{\beta_1}(\mathcal{G}_2) < \delta$ and $\mathfrak{p}_{\beta_2}(\mathcal{G}_1) < \delta$

imply an existence of an inverse temperature $\beta_t \in (\beta_1, \beta_2)$ and of two distinct B -shift ergodic Gibbs measures $\mu_1, \mu_2 \in \mathfrak{G}_{\beta_t}$ such that

$$\rho_{\mu_j}(\mathcal{G}_j) \geq 1 - \epsilon, \quad j = 1, 2. \quad (2.17)$$

The above assumptions (1) and (2) appear in some form in all existing proofs based on chessboard estimates; see Sect. 4 for some explicit examples. The conclusions about the set of coexistence points can be significantly strengthened when, on the basis of thermodynamic arguments and/or stochastic domination, the expected amount of goodness \mathcal{G}_2 increases (and \mathcal{G}_1 decreases) with increasing β . For $\epsilon \ll 1$ the phase diagram then features a unique (massive) jump at some β_t from states dominated by \mathcal{G}_1 to those dominated by \mathcal{G}_2 . Theorem 2.5 implies that the bulk of the values inside the jump are not found in any ergodic Gibbs state. Both Theorem 2.5 and Corollary 2.6 are proved in Sect. 3.2.

3. PROOFS OF MAIN RESULTS

We will assume that there is an ergodic Gibbs measure $\mu \in \mathfrak{G}_\beta$ that violates one of the conditions (2.15–2.16), and derive a contradiction. Various steps of the proof will be encapsulated in technical lemmas in Sect. 3.1; the actual proofs come in Sect. 3.2.

3.1 Technical lemmas.

Our first step is to convert the information about infinite-volume densities into a finite volume event. Using the sites from Λ_{N-1} to translate the B -block Λ_B by multiples of B in each coordinate direction, we get $\bigcup_{x \in \Lambda_{N-1}} (\Lambda_B + Bx) = \Lambda_{NB}$. Similarly, considering translates of Λ_{NB} by vectors NBx where $x \in \Lambda_{M-1}$, we get $\bigcup_{x \in \Lambda_{M-1}} (\Lambda_{NB} + NBx) = \Lambda_{MNB}$. The important point is that, while the neighboring translates $\Lambda_{NB} + NBx$ and $\Lambda_{NB} + NBx + NBx$ are not disjoint, they have only one of their $(d-1)$ -dimensional sides in common.

Let \mathcal{B}_N and $\mathcal{E}_{j,N}$, $j = 1, \dots, r$, be events defined by

$$\mathcal{B}_N = \{R_N(\mathcal{B}) > \epsilon\} \quad (3.1)$$

and

$$\mathcal{E}_{j,N} = \{R_N(\mathcal{G}_j) > \epsilon\}, \quad j = 1, \dots, r. \quad (3.2)$$

Introducing the event

$$\mathcal{E}_N = \mathcal{B}_N \cup \bigcup_{1 \leq i < j \leq r} (\mathcal{E}_{i,N} \cap \mathcal{E}_{j,N}) \quad (3.3)$$

and the fraction $R_{M,N}(\mathcal{E}_N)$ of BN -blocks (in Λ_{MNB}) in which \mathcal{E}_N occurs,

$$R_{M,N}(\mathcal{E}_N) = \frac{1}{|\Lambda_{M-1}|} \sum_{x \in \Lambda_{M-1}} \mathbf{1}_{\mathcal{E}_N \circ \tau_{NBx}}, \quad (3.4)$$

we have:

Lemma 3.1 *Let $\epsilon < 1/2$ and consider a B -shift ergodic Gibbs measure $\mu \in \mathfrak{G}_\beta$ that violates one of the conditions (2.15–2.16). Then there exists an $N_0 < \infty$ and, for each $N \geq N_0$, there exists an $M_0 = M_0(N)$ such that for all $N \geq N_0$ and all $M \geq M_0(N)$, one has*

$$\mu(R_{M,N}(\mathcal{E}_N) > 1/2) > \frac{1}{2N^d}. \quad (3.5)$$

Proof. The proof is based on a two-fold application of the Pointwise Ergodic Theorem. Indeed, by ergodicity of μ and Fatou's lemma we know that

$$\liminf_{N \rightarrow \infty} \mu(\mathcal{B}_N) \geq \mu(\rho_\mu(\mathcal{B}) > \epsilon) \quad (3.6)$$

and

$$\liminf_{N \rightarrow \infty} \mu(\mathcal{E}_{i,N} \cap \mathcal{E}_{j,N}) \geq \mu(\{\rho_\mu(\mathcal{G}_i) > \epsilon\} \cap \{\rho_\mu(\mathcal{G}_j) > \epsilon\}). \quad (3.7)$$

But μ violates one of the conditions (2.15–2.16) and so either $\rho_\mu(\mathcal{B}) > \epsilon$ or $\rho_\mu(\mathcal{G}_i) > \epsilon$ and $\rho_\mu(\mathcal{G}_j) > \epsilon$ for some $i \neq j$. All of these inequalities are valid μ -almost surely and so it follows that

$$\mu(\mathcal{E}_N) \xrightarrow{N \rightarrow \infty} 1. \quad (3.8)$$

Now, let us fix N so that $\mu(\mathcal{E}_N) \geq 3/4$. Then ergodicity with respect to translates by multiples of B implies that

$$\begin{aligned} \mu\left(\bigcup_{y \in \Lambda_{N-1}} \{R_{M,N}(\mathcal{E}_N) \circ \tau_{By} > 1/2\}\right) &\geq \mu\left(\frac{1}{N^d} \sum_{y \in \Lambda_{N-1}} R_{M,N}(\mathcal{E}_N) \circ \tau_{By} > \frac{1}{2}\right) \\ &= \mu(R_{MN}(\mathcal{E}_N) > 1/2) \xrightarrow{M \rightarrow \infty} 1. \end{aligned} \quad (3.9)$$

It follows that the left-hand side exceeds $1/2$ once M is sufficiently large, which in conjunction with subadditivity and τ_{By} -invariance of μ directly implies (3.5). \square

Our next task will be to express \mathcal{E}_N solely in terms of conditions on bad B -blocks in $\Lambda_{NB} = \bigcup_{x \in \Lambda_{N-1}} (\Lambda_B + Bx)$. Given two distinct sites $x, y \in \Lambda_{N-1}$, let $\{x \leftrightarrow y\}$ denote the event that there is no nearest-neighbor path $\pi = (x_1, \dots, x_k)$ on Λ_{N-1} such that

- (1) π connects x to y , i.e., $x_1 = x$ and $x_k = y$.
- (2) all B -blocks “along” π are good, i.e., $\tau_{Bx_j}(\mathcal{B}^c)$ occurs for all $j = 1, \dots, k$.

Note that $\{x \leftrightarrow y\}$ automatically holds when one of the blocks $\Lambda_B + Bx$ or $\Lambda_B + By$ is bad. Further, let Y_N be the ($\mathcal{F}_{\Lambda_{NB}}$ -measurable) random variable

$$Y_N = \#\{(x, y) \in \Lambda_{N-1} \times \Lambda_{N-1} : x \neq y \text{ \& } x \leftrightarrow y\} \quad (3.10)$$

and let \mathcal{C}_N be the event

$$\mathcal{C}_N = \{Y_N \geq (\epsilon N^d)^2\}. \quad (3.11)$$

Conditions (1) and (2) from Theorem 2.5 now directly imply:

Lemma 3.2 *For all N , we have $\mathcal{E}_N \subset \mathcal{C}_N$.*

Proof. Clearly, we have $\mathcal{B}_N \subset \mathcal{C}_N$, and so we only have to show that

$$\mathcal{E}_{i,N} \cap \mathcal{E}_{j,N} \subset \mathcal{C}_N, \quad 1 \leq i < j \leq r. \quad (3.12)$$

Let us fix $i \neq j$ and recall that on $\mathcal{E}_{i,N} \cap \mathcal{E}_{j,N}$, at least an ϵ -fraction of all B -blocks in Λ_{NB} will be i -good and at least an ϵ -fraction of them will be j -good. By conditions (1) and (2) from Theorem 2.5, no two B -blocks of different type of goodness can be connected by a path of good B -blocks, and so there are at least $(\epsilon N^d)^2$ pairs of distinct B -blocks in Λ_{NB} that are not connected to each other by a path of good blocks. This is exactly what defines the event \mathcal{C}_N . \square

The events \mathcal{E}_N and \mathcal{C}_N have the natural interpretation as NB -block events on \mathbb{T}_L whenever L is divisible by NB . If \mathcal{A} is such an NB -block event, let $\tilde{\mathfrak{p}}_\beta(\mathcal{A})$ denote the analogue of the quantity from (2.11) where the θ_t 's now involve translations by multiples of NB . Our next technical lemma provides an estimate on $\tilde{\mathfrak{p}}_\beta(\mathcal{C}_N)$ in terms of $\mathfrak{p}_\beta(\mathcal{B})$:

Lemma 3.3 *Let d be the dimension of the underlying lattice and suppose that $d \geq 2$. For each $\epsilon > 0$ —underlying the definitions of \mathcal{B}_N , \mathcal{E}_N and \mathcal{C}_N —and each $\eta > 0$, there exists a number $\delta = \delta(\epsilon, \eta, d) > 0$ such that if $\mathfrak{p}_\beta(\mathcal{B}) < \delta$, then $\tilde{\mathfrak{p}}_\beta(\mathcal{C}_N) < \eta$.*

Proof. Let us use $\Pi_{L,\beta}(\mathcal{C}_N)$ to abbreviate the quantity

$$\Pi_{L,\beta}(\mathcal{C}_N) = \mathbb{P}_{L,\beta} \left(\bigcap_{t \in \tilde{\mathbb{T}}} \theta_t(\mathcal{C}_N) \right), \quad (3.13)$$

where $\tilde{\mathbb{T}} = \mathbb{T}_{L/(NB)}$ is the factor torus in the present context. Observing that \mathcal{C}_N is preserved by reflections through the “midplanes” of Λ_{NB} , a multivariate version of Chebyshev’s inequality then yields

$$\Pi_{L,\beta}(\mathcal{C}_N) \leq \mathbb{E}_{L,\beta} \left(\prod_{t \in \tilde{\mathbb{T}}} \frac{Y_N \circ \tau_{BNt}}{(\epsilon N^d)^2} \right). \quad (3.14)$$

Here $\mathbb{E}_{L,\beta}$ is the expectation with respect to $\mathbb{P}_{L,\beta}$.

To estimate the right-hand side of (3.14), we will rewrite Y_N as a sum. Let $x, y \in \Lambda_{N-1}$ be distinct. A connected subset $\Gamma \subset \Lambda_{N-1}$ is said to *separate x from y* (in Λ_{N-1}) if each nearest-neighbor path π from x to y on Λ_{N-1} intersects Γ . We use $\mathfrak{S}(x, y)$ to denote the set of all such sets $\Gamma \subset \Lambda_{N-1}$. Notice that $\{x\}, \{y\} \in \mathfrak{S}(x, y)$. We claim that, whenever (x, y) is a pair of points contributing to Y_N , there exists $\Gamma \in \mathfrak{S}(x, y)$ separating x from y such that every block $\Lambda_B + Bz$ with $z \in \Gamma$ is bad. Indeed, if $\Lambda_B + Bx$ is a bad block we take $\Gamma = \{x\}$. If $\Lambda_B + Bx$ is a good block, then we define \mathcal{C}_x to be the maximal connected subset of Λ_{N-1} containing x such that $\Lambda_B + Bz$ is a good block for all $z \in \mathcal{C}_x$, and let Γ be its external boundary. Using $\mathbf{1}_\Gamma$ to

denote the indicator of the event that every block $\Lambda_B + Bz$ with $z \in \Gamma$ is bad, we get

$$Y_N \leq \sum_{x,y \in \Lambda_{N-1}} \sum_{\Gamma \in \mathfrak{S}(x,y)} \mathbf{1}_\Gamma. \quad (3.15)$$

Let $K = (\frac{L}{BN})^d$ be the volume of the factor torus and let $\mathbf{t}_1, \dots, \mathbf{t}_K$ be an ordering of all sites of $\tilde{\mathbb{T}}$. Then we have

$$\Pi_{L,\beta}(\mathcal{C}_N) \leq \frac{1}{(\epsilon N^d)^{2K}} \sum_{\substack{(x_j, y_j) \\ j=1, \dots, K}} \sum_{\Gamma_1, \dots, \Gamma_K} \mathbb{E}_{L,\beta} \left(\prod_{j=1}^K \mathbf{1}_{\Gamma_j \circ \tau_{BN\mathbf{t}_j}} \right), \quad (3.16)$$

where the first sum runs over collections of pairs (x_j, y_j) , $j = 1, \dots, K$, of distinct sites in Λ_{N-1} and the second sum is over all collections of separating surfaces $\Gamma_j \in \mathfrak{S}(x_j, y_j)$, $j = 1, \dots, K$.

To estimate the right-hand side of (3.16) we define $\mathfrak{p}_{L,\beta}(\mathcal{B})$ to be the quantity on the right-hand side of (2.11), before taking the limit $L \rightarrow \infty$, with $\mathcal{A} = \mathcal{B}$. Since each indicator $\mathbf{1}_{\Gamma_j \circ \tau_{BN\mathbf{t}_j}}$ enforces bad blocks $\Lambda_B + B(z + N\mathbf{t}_j)$ for $z \in \Gamma_j$, and the set of blocks $\Lambda_B + B(z + N\mathbf{t}_j)$, $z \in \Lambda_{N-1}$, is, for $\mathbf{t}_i \neq \mathbf{t}_j$, disjoint from the set $\Lambda_B + B(z + N\mathbf{t}_i)$, $z \in \Lambda_{N-1}$, we can use chessboard estimates (Theorem 2.4) to get

$$\mathbb{E}_{L,\beta} \left(\prod_{j=1}^K \mathbf{1}_{\Gamma_j \circ \tau_{BN\mathbf{t}_j}} \right) \leq [\mathfrak{p}_{L,\beta}(\mathcal{B})]^{|\Gamma_1| + \dots + |\Gamma_K|}. \quad (3.17)$$

A standard contour-counting argument now shows that, for any distinct $x, y \in \Lambda_{N-1}$,

$$\sum_{\Gamma \in \mathfrak{S}(x,y)} [\mathfrak{p}_{L,\beta}(\mathcal{B})]^{|\Gamma|} \leq c_1 \mathfrak{p}_{L,\beta}(\mathcal{B})^d \quad (3.18)$$

with some constant $c_1 = c_1(d)$, provided that $\mathfrak{p}_{L,\beta}(\mathcal{B})$ is sufficiently small. The sum over collections of pairs (x_j, y_j) , $j = 1, \dots, K$, contains at most $(N^{2d})^K$ terms, allowing us to bound

$$\Pi_{L,\beta}(\mathcal{C}_N) \leq \left(\frac{c_1 \mathfrak{p}_{L,\beta}(\mathcal{B})^d}{\epsilon^2} \right)^K. \quad (3.19)$$

Since $\Pi_{L,\beta}(\mathcal{C}_N)^{1/K} \rightarrow \tilde{\mathfrak{p}}_\beta(\mathcal{C}_N)$ as $L \rightarrow \infty$, it follows that $\tilde{\mathfrak{p}}_\beta(\mathcal{C}_N) \leq c_1 \mathfrak{p}_\beta(\mathcal{B})^d \epsilon^{-2}$, which for $\mathfrak{p}_\beta(\mathcal{B})$ small enough, can be made smaller than any η initially prescribed. \square

Our final technical ingredient is an estimate on the Radon-Nikodym derivative of a Gibbs measure $\mu \in \mathfrak{G}_\beta$ and the torus measure at the same temperature:

Lemma 3.4 *Let $\Lambda_L \subset \mathbb{Z}^d$ be an L -block and let \mathbb{T}_{2L} be a torus of side $2L$. Let us view Λ_L as embedded into \mathbb{T}_{2L} and let $\mathbb{P}_{2L,\beta}$ be the torus Gibbs measure on \mathbb{T}_{2L} . Then for any $a > 0$ there exists L_0 such that*

$$e^{-\beta a L^d} \mathbb{P}_{2L,\beta}(\mathcal{A}) \leq \mu(\mathcal{A}) \leq e^{\beta a L^d} \mathbb{P}_{2L,\beta}(\mathcal{A}). \quad (3.20)$$

for all $L \geq L_0$, any $\mu \in \mathfrak{G}_\beta$, and any \mathcal{F}_{Λ_L} -measurable event \mathcal{A} .

Proof. For finite-range interaction, this lemma is completely standard. However, since our setting includes also interactions with infinite range, we provide a complete proof. We will prove only the right-hand side of the above inequality; the other side is completely analogous.

First, from the DLR equation we know that there exists a configuration $s = (s_x)_{x \in \mathbb{Z}^d}$, such that

$$\mu(\mathcal{A} | \mathcal{F}_{\Lambda^c})(s) \geq \mu(\mathcal{A}) \quad (3.21)$$

with the left-hand side of the form (2.3). Let s' be a configuration on \mathbb{T}_{2L} . We will show that $\mu(\cdot | \mathcal{F}_{\Lambda_L^c})(s)$ and $\mathbb{P}_{2L, \beta}(\cdot | \mathcal{F}_{\Lambda_L^c})(s')$ are absolutely continuous with respect to each other—as measures on \mathcal{F}_{Λ_L} —and the Radon-Nikodym derivative is bounded above by $e^{\beta a L^d}$ regardless of the “boundary conditions” s and s' .

Suppose that $s'_x = s_x$ for all $x \in \Lambda_L$ and let \bar{s}' be its $2L$ -periodic extension to all of \mathbb{Z}^d . Then the Radon-Nikodym derivative of $\mathbb{P}_{2L, \beta}(\cdot | \mathcal{F}_{\Lambda_L^c})(s')$ with respect to the product measure $\prod_{x \in \Lambda_L} \nu_0(ds_x)$ is $e^{-\beta H_{\Lambda_L}(\bar{s}')}/Z_{\Lambda_L}(\bar{s}'_{\Lambda_L^c})$ while that of $\mu(\cdot | \mathcal{F}_{\Lambda_L^c})(s)$ is $e^{-\beta H_{\Lambda_L}(s)}/Z_{\Lambda_L}(s_{\Lambda_L^c})$. It thus suffices to show, uniformly in $(s_x)_{x \in \Lambda_L}$, that

$$|H_{\Lambda_L}(s) - H_{\Lambda_L}(\bar{s}')| \leq \frac{a}{2} L^d \quad (3.22)$$

once L is sufficiently large. To this end, we first note that

$$|H_{\Lambda_L}(s) - H_{\Lambda_L}(\bar{s}')| \leq 2 \sum_{\substack{A: A \cap \Lambda_L \neq \emptyset \\ A \cap \Lambda_L^c \neq \emptyset}} \|\Phi_A\|_{\infty}. \quad (3.23)$$

To estimate the right-hand side, we will decompose Λ_L into “shells,” $\Lambda_n \setminus \Lambda_{n-1}$, and use the fact that if A intersects $\Lambda_n \setminus \Lambda_{n-1}$ as well as Λ_L^c , then the diameter of A must be at least $L - n$. Using the translation invariance of the interactions, we thus get

$$\sum_{\substack{A: A \cap \Lambda_L \neq \emptyset \\ A \cap \Lambda_L^c \neq \emptyset}} \|\Phi_A\|_{\infty} \leq \sum_{n=1}^L |\Lambda_n \setminus \Lambda_{n-1}| \sum_{\substack{A: 0 \in A \\ \text{diam}(A) \geq L-n}} \|\Phi_A\|_{\infty}. \quad (3.24)$$

But $\|\Phi\| < \infty$ implies that the second sum tends to zero as $L - n \rightarrow \infty$ and since $|\Lambda_n \setminus \Lambda_{n-1}| = o(L^d)$ while $\sum_{1 \leq n \leq L} |\Lambda_n \setminus \Lambda_{n-1}| = L^d$, the result is thus $o(L^d)$. In particular, for L sufficiently large, the right-hand side of (3.23) will be less than $\frac{a}{2} L^d$. \square

3.2 Proofs of Theorem 2.5 and Corollary 2.6.

Now we are ready to prove our main theorem:

Proof of Theorem 2.5. Fix $\epsilon < 1/2$ and let $\mu \in \mathfrak{G}_{\beta}$ be a B -shift ergodic Gibbs measure for which one of the conditions (2.15–2.16) fails. Applying Lemma 3.1 and the inclusion in Lemma 3.2 we find that

$$\mu(R_{M, N}(C_N) > 1/2) > \frac{1}{2N^d} \quad (3.25)$$

once $N \geq N_0$ and $M \geq M_0(N)$. Now, consider the torus \mathbb{T}_L of side $L = 2MNB$ and embed $\Lambda_{MNB} = \bigcup_{x \in \Lambda_{M-1}} (\Lambda_{NB} + NBx)$ into \mathbb{T}_L in the “usual” way. By Lemma 3.4 we know that for any fixed $N \geq N_0$, there exists a sequence a_M of positive numbers with $a_M \downarrow 0$ as $M \rightarrow \infty$, such that we have

$$\mathbb{P}_{L, \beta}(R_{M, N}(C_N) > 1/2) > \frac{1}{2N^d} e^{-\beta(NB)^d a_M M^d}, \quad M \rightarrow \infty. \quad (3.26)$$

Our goal is to show that, once N is chosen sufficiently large, the left-hand side is exponentially small in M^d , thus arriving at a contradiction.

By conditioning on which of the $M^d/2$ translates of Λ_{BN} have \mathcal{C}_N satisfied, and applying the chessboard estimates in blocks of side NB , we get

$$\mathbb{P}_{L,\beta}(\mathcal{R}_{M,N}(\mathcal{C}_N) > 1/2) \leq 2^{M^d} \tilde{\mathfrak{p}}_{2L,\beta}(\mathcal{C}_N)^{M^d/2}, \quad (3.27)$$

where $\tilde{\mathfrak{p}}_{2L,\beta}(\mathcal{C}_N)$ is the finite-torus version of $\tilde{\mathfrak{p}}_\beta(\mathcal{C}_N)$. Next we choose $\eta < 1/4$ and let $\delta > 0$ and $N \geq N_0$ be such that the bounds in Lemma 3.3 apply. Then for all sufficiently large M (and hence all large L) we have $\tilde{\mathfrak{p}}_{2L,\beta}(\mathcal{C}_N) < \eta$ and so

$$\mathbb{P}_{L,\beta}(\mathcal{R}_{M,N}(\mathcal{C}_N) > 1/2) \leq (4\eta)^{M^d/2}. \quad (3.28)$$

But this is true for all $M \gg 1$ and so the bound (3.26) must be false. Hence, no such $\mu \in \mathfrak{G}_\beta$ could exist to begin with; i.e., (2.15–2.16) must hold for all B -shift ergodic $\mu \in \mathfrak{G}_\beta$. \square

To finish our proofs, we will also need to establish our claims concerning phase coexistence:

Proof of Corollary 2.6. Suppose that ϵ and δ are such that Theorem 2.5 applies. By condition (1), the conclusions (2.15–2.16) of this theorem are thus available for all $\beta \in [\beta_1, \beta_2]$. This implies

$$\rho_\mu(\mathcal{G}_j) \in [0, \epsilon] \cup [1 - \epsilon, 1], \quad j = 1, 2, \quad (3.29)$$

for every B -shift ergodic $\mu \in \mathfrak{G}_\beta$ at every $\beta \in [\beta_1, \beta_2]$. We claim that $\rho_\mu(\mathcal{G}_2)$ is small in every ergodic state $\mu \in \mathfrak{G}_{\beta_1}$. Indeed, by Lemma 6.3 of [4] and condition (2) of the corollary, we have

$$\mathfrak{p}_{\beta_1}(\mathcal{B} \cup \mathcal{G}_2) \leq \mathfrak{p}_{\beta_1}(\mathcal{B}) + \mathfrak{p}_{\beta_1}(\mathcal{G}_2) < 2\delta. \quad (3.30)$$

Hence, if the δ in Corollary 2.6 was so small that Theorem 2.5 applies for some $\epsilon < 1/2$ even when δ is replaced by 2δ , we can regard $\mathcal{B} \cup \mathcal{G}_2$ as a bad event at $\beta = \beta_1$ and conclude that $\rho_\mu(\mathcal{G}_2) < 1/2$, and hence $\rho_\mu(\mathcal{G}_2) \leq \epsilon$, by (3.29), in every ergodic $\mu \in \mathfrak{G}_{\beta_1}$. A similar argument proves that $\rho_\mu(\mathcal{G}_1) \leq \epsilon$ in every ergodic $\mu \in \mathfrak{G}_{\beta_2}$. Usual weak-limit arguments then yield the existence of at least one point $\beta_t \in (\beta_1, \beta_2)$ where both types of goodness coexist. \square

4. APPLICATIONS

The formulation of our main result is somewhat abstract. In the present section, we will pick several models in which phase coexistence has been proved using chessboard estimates and use them to demonstrate the consequences of our main theorem. Although we will try to stay rather brief, we will show that, generally, the hypothesis of our main result—i.e., the assumption on smallness of the parameter $\mathfrak{p}_\beta(\mathcal{B})$ —is directly implied by the calculations already carried out in the corresponding papers. The reader should consult the original articles for more motivation and further details concerning the particular models.

4.1 Potts model.

The q -state Potts model serves as a paradigm of order-disorder transitions. The existence of the transition has been proved by chessboard estimates in [16]. While the completeness of the phase diagram has, in the meantime, been established with the help of Pirogov-Sinai theory [18], we

find it useful to illustrate our general claims on this rather straightforward example. Later on we will pass to more complex systems where no form of “completeness” has so far been proved.

The spins σ_x of the q -state Potts model take values in the set $\{1, \dots, q\}$ with *a priori* equal probabilities. The formal Hamiltonian is

$$H(\sigma) = - \sum_{\langle x, y \rangle} \delta_{\sigma_x, \sigma_y}, \quad (4.1)$$

where $\langle x, y \rangle$ runs over all (unordered) nearest-neighbor pairs in \mathbb{Z}^d . The states of minimal energy have all neighboring spins equal, and so we expect that low temperature states are dominated by nearly constant spin-configurations. On the other hand, at high temperatures the spins should be nearly independent and, in particular, neighboring spins will typically be different from each other. This leads us to consider the following good events on 1-block Λ_1 :

$$\begin{aligned} \mathcal{G}^{\text{dis}} &= \{\sigma : \sigma_x \neq \sigma_y \text{ for all } x, y \in \Lambda_1, |x - y| = 1\}, \\ \mathcal{G}^{\text{ord}, m} &= \{\sigma : \sigma_x = m \text{ for all } x \in \Lambda_1\}, \quad m = 1, \dots, q. \end{aligned} \quad (4.2)$$

Using similar events, it was proved [16] that, for $d \geq 2$ and q sufficiently large, there exists an inverse temperature β_t and $q + 1$ ergodic Gibbs states $\mu^{\text{dis}} \in \mathfrak{G}_{\beta_t}$ and $\mu^{\text{ord}, m} \in \mathfrak{G}_{\beta_t}$, $m = 1, \dots, q$, such that the corresponding 1-block densities satisfy

$$\rho_{\mu^{\text{dis}}}(\mathcal{G}^{\text{dis}}) \geq 1 - \epsilon \quad (4.3)$$

and

$$\rho_{\mu^{\text{ord}, m}}(\mathcal{G}^{\text{ord}, m}) \geq 1 - \epsilon, \quad m = 1, \dots, q, \quad (4.4)$$

where $\epsilon = \epsilon(q)$ tends to zero as $q \rightarrow \infty$. In addition, monotonicity of the energy density as a function of β can be invoked to show that $\rho_{\mu}(\mathcal{G}^{\text{dis}})$ is large in all translation-invariant $\mu \in \mathfrak{G}_{\beta}$ when $\beta < \beta_t$, while it is small in all such states when $\beta > \beta_t$.

The full completeness [18] asserts that above mentioned $q + 1$ states exhaust the set of all shift-ergodic Gibbs states in \mathfrak{G}_{β_t} . A weaker claim follows as a straightforward application of our Theorem 2.5: There is *no* shift-ergodic Gibbs state $\mu \in \mathfrak{G}_{\beta_t}$ such that

$$\rho_{\mu}(\mathcal{G}^{\text{dis}}) \in [\epsilon, 1 - \epsilon] \quad \text{or} \quad \rho_{\mu}(\mathcal{G}^{\text{ord}, m}) \in [\epsilon, 1 - \epsilon], \quad (4.5)$$

for some $m = 1, \dots, q$. The main hypothesis of our theorem amounts to the smallness of the quantity $\mathfrak{p}_{\beta}(\mathcal{B})$, where

$$\mathcal{B} = \left(\mathcal{G}^{\text{dis}} \cup \bigcup_{m=1}^q \mathcal{G}^{\text{ord}, m} \right)^c, \quad (4.6)$$

which in turn boils down to an estimate on the probability of the disseminated event \mathcal{B} on the right-hand side of (2.11). The needed estimate coincides with the bound provided in [16] by evaluating directly (i.e., “by hand”) the energy and the number of contributing configurations. The result—which in [16] appears right before the last formula on p. 506 is used to produce (4.4)—reads

$$\mathfrak{p}_{\beta}(\mathcal{B}) \leq \left[\frac{q^{d-2-(d-1)}}{(q-2d)^d} \right]^{\frac{1}{2d}}. \quad (4.7)$$

This implies the needed bound once $q \gg 1$.

Remark 4.1 Analogous calculations establish the corresponding forbidden gap in more complicated variants of the Potts model; see e.g. [3].

4.2 Intermediate phases in dilute spin systems.

Another instance where our results provide new insight are dilute annealed ferromagnets exhibiting staggered order phases at intermediate temperatures. These systems have been studied in the context of both discrete [6] and continuous spins [7]. The characteristic examples of these classes are the *site-diluted Potts model* with the Hamiltonian

$$H(n, \sigma) = - \sum_{\langle x, y \rangle} n_x n_y (\delta_{\sigma_x, \sigma_y} - 1) - \lambda \sum_x n_x - \kappa \sum_{\langle x, y \rangle} n_x n_y \quad (4.8)$$

and the *site-diluted XY-model* with the Hamiltonian

$$H(n, \phi) = - \sum_{\langle x, y \rangle} n_x n_y [\cos(\phi_x - \phi_y) - 1] - \lambda \sum_x n_x - \kappa \sum_{\langle x, y \rangle} n_x n_y. \quad (4.9)$$

Here, as before, $\sigma_x \in \{1, \dots, q\}$ are the Potts spins, $\phi_x \in [-\pi, \pi)$ are variables representing the “angle” of the corresponding $O(2)$ -spins, and $n_x \in \{0, 1\}$ indicates the presence or absence of a particle (that carries the Potts spin σ_x or the angle variable ϕ_x) at site x .

On the basis of “usual” arguments, the high temperature region is characterized by disordered configurations while the low temperatures features configurations with a strong (local) order, at least at small-to-intermediate dilutions. The phenomenon discovered in [6, 7] is the existence of a region of intermediate temperatures and chemical potentials, sandwiched between the low temperature/high density ordered region and the high temperature/low density disordered region, where typical configurations exhibit preferential occupation of one of the even/odd sublattices. The appearance of such states is due to an *effective entropic repulsion*. Indeed, at low temperatures the spins on particles at neighboring sites are forced to be (nearly) aligned while if a particle is completely isolated, its spin is permitted to enjoy the full freedom of the available spin space. Hence, at intermediate temperatures and moderate dilutions, there is an entropic advantage for the particles to occupy only one of the sublattices.

Let us concentrate on the portion of the phase boundary between the staggered region and the low temperature region. The claim can be stated uniformly for both systems in (4.8–4.9) provided we introduce the relevant good events in terms of occupation variable n . Namely, we let:

$$\begin{aligned} \mathcal{G}^{\text{dense}} &= \{(\sigma, n) : n_x = 1 \text{ for all } x \in \Lambda_1\}, \\ \mathcal{G}^{\text{even}} &= \{(\sigma, n) : n_x = \mathbf{1}_{\{x \text{ even}\}} \text{ for all } x \in \Lambda_1\}, \\ \mathcal{G}^{\text{odd}} &= \{(\sigma, n) : n_x = \mathbf{1}_{\{x \text{ odd}\}} \text{ for all } x \in \Lambda_1\}. \end{aligned} \quad (4.10)$$

Again, using slightly modified versions of these events, it was shown in [6, 7] that there exist positive numbers $\epsilon, \kappa_0 \ll 1$ and, for every $\kappa \in (0, \kappa_0)$, an interval $I(\kappa) \subset \mathbb{R}$ such that the following is true: For any $\lambda \in I$ there exist inverse temperatures $\beta_1(\kappa, \lambda)$ and $\beta_2(\kappa, \lambda)$, and a transition temperature $\beta_t(\kappa, \lambda) \in [\beta_1, \beta_2]$ such that

- (1) for any $\beta \in [\beta_t, \beta_2]$ there exists an “densely occupied” state $\mu^{\text{dense}} \in \mathfrak{G}_\beta$, for which

$$\rho_{\mu^{\text{dense}}}(\mathcal{G}^{\text{dense}}) \geq 1 - \epsilon, \quad (4.11)$$

(2) for any $\beta \in [\beta_1, \beta_t]$ there exist two states $\mu^{\text{even}}, \mu^{\text{odd}} \in \mathfrak{G}_\beta$ satisfying

$$\rho_{\mu^{\text{even}}}(\mathcal{G}^{\text{even}}) \geq 1 - \epsilon \quad \text{and} \quad \rho_{\mu^{\text{odd}}}(\mathcal{G}^{\text{odd}}) \geq 1 - \epsilon. \quad (4.12)$$

The error ϵ is of order $\beta^{-1/8}$ (cf. the bound (2.15) in [7]) in the case of the XY-model in $d = 2$, and it tends zero as $q \rightarrow \infty$ in the case of the diluted Potts model.

A somewhat stronger conclusion can be made for the diluted Potts model. Namely, at $\beta = \beta_t$, there are actually $q + 2$ distinct states, two staggered states μ^{even} and μ^{odd} and q ordered states $\mu^{\text{dense}, m}$, with the latter characterized by the condition

$$\rho_{\mu^{\text{dense}, m}}(\mathcal{G}^{\text{dense}, m}) \geq 1 - \epsilon, \quad (4.13)$$

where

$$\mathcal{G}^{\text{dense}, m} = \{(\sigma, n) : n_x = 1 \text{ and } \sigma_x = m \text{ for all } x \in \Lambda_1\}. \quad (4.14)$$

It is plausible that an analogous conclusion applies to the XY-model in $d \geq 3$ because there the low-temperature phase should exhibit magnetic order. However, in $d = 2$ such long-range order is not permitted by the Mermin-Wagner theorem and so there one expects to have only 3 distinct ergodic Gibbs states at β_t .

A weaker form of the expected conclusions is an easy consequence of our Theorem 2.5: There exists no shift-ergodic Gibbs state $\mu \in \mathfrak{G}_{\beta_t}$ such that

$$\rho_\mu(\mathcal{G}) \in [\epsilon, 1 - \epsilon] \quad \text{for some } \mathcal{G} \in \{\mathcal{G}^{\text{dense}}, \mathcal{G}^{\text{even}}, \mathcal{G}^{\text{odd}}\} \quad (4.15)$$

and, in the case of diluted Potts model, also

$$\rho_\mu(\mathcal{G}^{\text{dense}, m}) \in [\epsilon, 1 - \epsilon] \quad \text{for some } m \in \{1, \dots, q\}. \quad (4.16)$$

In particular, no ergodic Gibbs state $\mu \in \mathfrak{G}_{\beta_t}$ has particle density in $[\epsilon, 1/2 - \epsilon] \cup [1/2 + \epsilon, 1 - \epsilon]$. The proof of these observations goes by noting that the smallness of $\mathfrak{p}_\beta(\mathcal{B})$ for the bad event $\mathcal{B} = (\mathcal{G}^{\text{dense}} \cup \mathcal{G}^{\text{even}} \cup \mathcal{G}^{\text{odd}})^c$ is a direct consequence of the corresponding bounds from [6, 7] of the ‘‘contour events.’’ In the case of the XY-model in dimension $d = 2$, this amounts to the bounds (2.9) and (2.15) from [7].

Remark 4.2 A more general class of models, with spin taking values in a Riemannian manifold, is also considered in [7]. A related phase transition in an annealed diluted $O(n)$ Heisenberg ferromagnet has been proved in [8].

4.3 Order-by-disorder transitions.

Another class of systems where our results provide new information are the $O(2)$ -nearest and next-nearest neighbor antiferromagnet [2], the 120-degree model [4], and the orbital-compass model [5]. All of these are continuum-spin systems whose common feature is that the infinite degeneracy of the ground states is broken, at positive temperatures, by long-wavelength (spin-wave) excitation. We will restrict our attention to the first of these models, the $O(2)$ -nearest and next-nearest neighbor antiferromagnet. The other two models are somewhat more complicated—particularly, due to the presence of non-translation invariant ground states—but the conclusions are fairly analogous.

Consider a spin system on \mathbb{Z}^2 whose spins, \mathbf{S}_x , take values on the unit circle in \mathbb{R}^2 with *a priori* uniform distribution. The Hamiltonian is

$$H(\mathbf{S}) = \sum_x (\mathbf{S}_x \cdot \mathbf{S}_{x+\hat{e}_1+\hat{e}_2} + \mathbf{S}_x \cdot \mathbf{S}_{x+\hat{e}_1-\hat{e}_2}) + \gamma \sum_x (\mathbf{S}_x \cdot \mathbf{S}_{x+\hat{e}_1} + \mathbf{S}_x \cdot \mathbf{S}_{x+\hat{e}_2}), \quad (4.17)$$

where \hat{e}_1 and \hat{e}_2 are the unit vectors in the coordinate lattice directions and the dot denotes the usual scalar product. Note that both nearest and next-nearest neighbors are coupled antiferromagnetically but with a different strength. The following are the ground state configurations for $\gamma \in (-2, 2)$: Both even and odd sublattices enjoy a Néel (antiferromagnetic) order, but the relative orientation of these sublattice states is arbitrary.

It is clear that, at low temperatures, the configurations will be locally near one of the aforementioned ground states. Due to the continuous nature of the spins, the fluctuation spectrum is dominated by “harmonic perturbations,” a.k.a. *spin waves*. A heuristic spin-wave calculation (cf. [4, Sect. 2.2] for an example in the context of the 120-degree model) suggests that among all 2π possible relative orientations of the sublattices, the parallel and the antiparallel orientations are those entropically most favorable. And, indeed, as was proved in [2], there exist two 2-periodic Gibbs states μ_1 and μ_2 with the corresponding type of long-range order. However, the existence of Gibbs states with other relative orientations has not been ruled out.

We will now state a stronger version of [2, Theorem 2.1]. Let B be a large even integer and consider two B -block events \mathcal{G}_1 and \mathcal{G}_2 defined as follows: fixing a positive $\kappa \ll 1$, let

$$\mathcal{G}_1 = \bigcap_{\substack{x, y \in \Lambda_B \\ (y-x) \cdot \hat{e}_2 = 0}} \{\mathbf{S}_x \cdot \mathbf{S}_y \geq 1 - \kappa\} \cap \bigcap_{x, x+\hat{e}_2 \in \Lambda_B} \{\mathbf{S}_x \cdot \mathbf{S}_{x+\hat{e}_2} \leq -1 + \kappa\}, \quad (4.18)$$

i.e., \mathcal{G}_1 enforces horizontal stripes all over Λ_B . The event \mathcal{G}_2 in turn enforces vertical stripes; the definition is as above with the roles of \hat{e}_1 and \hat{e}_2 interchanged. Then we have:

Theorem 4.3 *Let $\gamma \in (0, 2)$ and let $\kappa \ll 1$. For each $\epsilon > 0$ there exists a $\beta_0 \in (0, \infty)$ and, for each $\beta \geq \beta_0$, there exists an integer $B \geq 1$ such that for any $\mu \in \mathfrak{G}_\beta$ that is ergodic with respect to shifts by multiples of B we have*

$$\rho_\mu(\mathcal{G}_j) \in [0, \epsilon] \cup [1 - \epsilon, 1], \quad j = 1, 2. \quad (4.19)$$

In particular, there exist two ergodic Gibbs states $\mu_1, \mu_2 \in \mathfrak{G}_\beta$, such that

$$\rho_{\mu_j}(\mathcal{G}_j) \geq 1 - \epsilon, \quad j = 1, 2, \quad (4.20)$$

The second conclusion—the existence of Gibbs states with parallel and antiparallel relative orientation of the sublattices—was the main content of Theorem 2.1 of [2]. What we have added here is that the corresponding configurations dominate *all* ergodic Gibbs states. The $O(2)$ ground-state symmetry of the relative orientation of the sublattices is thus truly broken at positive temperatures, which bolsters significantly the main point of [2]. Note that no restrictions are posed on the overall orientation of the spins. Indeed, by the Mermin-Wagner theorem (see [2, Theorem 2.2]) every $\mu \in \mathfrak{G}_\beta$ is invariant under simultaneous rotations of all spins.

Proof of Theorem 4.3. As expected, the proof boils down to showing that, for a proper choice of scale B we have $\mathfrak{p}_\beta(\mathcal{B}) \ll 1$ for $\mathcal{B} = (\mathcal{G}_1 \cup \mathcal{G}_2)^c$. In [2] this is done by decomposing \mathcal{B} into more elementary events—depending on whether the “badness” comes from excessive energy or

insufficient entropy—and estimating each of them separately. The relevant bounds are proved in [2, Lemmas 4.4 and 4.5] and combined together in [2, Eq. (4.20)]. Applying Theorem 2.5 of the present paper, we thus know that every B -shift ergodic $\mu \in \mathfrak{G}_\beta$ is dominated either by blocks of type \mathcal{G}_1 or by blocks of type \mathcal{G}_2 . Since $\rho_\mu(\mathcal{B}) \leq \epsilon$ in all states, the existence of $\mu_1, \mu_2 \in \mathfrak{G}_\beta$ satisfying (4.20) follows by symmetry with respect to rotation (of the lattice) by 90-degrees. \square

4.4 Nonlinear vector models.

A class of models with continuous symmetry that are conceptually close to the Potts model has been studied recently by van Enter and Shlosman [11]. As for our previous examples with continuous spins, Pirogov-Sinai theory is not readily available and one has to rely on chessboard estimates. We will focus our attention on one example in this class, a *nonlinear ferromagnet*, although our conclusions apply with appropriate, and somewhat delicate, modifications also to liquid crystal models and lattice gauge models discussed in [11].

Let us consider an $O(2)$ -spin system on \mathbb{Z}^2 with spins parametrized by the angular variables $\phi_x \in (-\pi, \pi]$. The Hamiltonian is given by

$$H(\phi) = - \sum_{\langle x, y \rangle} \left(\frac{1 + \cos(\phi_x - \phi_y)}{2} \right)^p, \quad (4.21)$$

where p is a nonlinearity parameter. The *a priori* distribution of the ϕ_x 's is the Lebesgue measure on $(-\pi, \pi]$; the difference $\phi_x - \phi_y$ is always taken modulo 2π .

In order to define the good block events, we first split all bonds into three classes. Namely, given a configuration $(\phi_x)_{x \in \mathbb{Z}^2}$, we say that the bond $\langle x, y \rangle$ is

- (1) *strongly ordered* if $|\phi_x - \phi_y| \leq \frac{1}{C\sqrt{p}}$,
- (2) *weakly ordered* if $\frac{1}{C\sqrt{p}} < |\phi_x - \phi_y| < \frac{C}{\sqrt{p}}$, and
- (3) *disordered* if $|\phi_x - \phi_y| \geq \frac{C}{\sqrt{p}}$.

Here C is a large number to be determined later. If a bond is either strongly or weakly ordered, we will call it simply *ordered*.

On the basis of (4.21), it is clear that strictly ordered bonds are favored energetically while the disordered bonds are favored entropically. The main observation of [11] is that, at least in torus measures, ordered and disordered bonds are unlikely to occur in the same configuration. Unfortunately, ordered bonds can be continuously deformed to become disordered and *vice versa* and so to prove the existence of an order-disorder transition, one needs to supplement this by the observation that it is unlikely to have many bonds in the “borderline” region $|\phi_x - \phi_y| \approx \frac{C}{\sqrt{p}}$. In order to correct for this problem, and to enable the use of our main theorems, we will have to prove the existence of a free-energy barrier between the *strongly* ordered and disordered phases.

Let Λ_1 be a 1-block (i.e., a plaquette) and let us consider the following good events on Λ_1 : The event that all bonds on Λ_1 are strongly ordered,

$$\mathcal{G}_{\text{so}} = \left\{ |\phi_x - \phi_y| \leq \frac{1}{C\sqrt{p}} : \forall x, y \in \Lambda_1, |x - y| = 1 \right\}, \quad (4.22)$$

and the event that all bonds on Λ_1 are disordered,

$$\mathcal{G}_{\text{dis}} = \left\{ |\phi_x - \phi_y| \geq \frac{C}{\sqrt{p}} : \forall x, y \in \Lambda_1, |x - y| = 1 \right\}. \quad (4.23)$$

As usual, let $\mathcal{B} = (\mathcal{G}_{\text{so}} \cup \mathcal{G}_{\text{dis}})^c$ be the corresponding bad block event. Then we have:

Theorem 4.4 *For each $\epsilon > 0$ and each sufficiently large $C > 1$, there exists $p_0 > 0$ such that for all $p > p_0$, all $\beta \geq 0$ and all shift-ergodic $\mu \in \mathfrak{G}_\beta$, we have*

$$\rho_\mu(\mathcal{G}_{\text{dis}}), \rho_\mu(\mathcal{G}_{\text{so}}) \in [0, \epsilon] \cup [1 - \epsilon, 1] \quad (4.24)$$

and

$$\rho_\mu(\mathcal{B}) \leq \epsilon. \quad (4.25)$$

Moreover, for every $p > p_0$ there exists a number $\beta_t \in (0, \infty)$ and two distinct, shift-ergodic Gibbs states $\mu^{\text{so}}, \mu^{\text{dis}} \in \mathfrak{G}_{\beta_t}$ such that

$$\rho_{\mu^{\text{so}}}(\mathcal{G}_{\text{so}}) \geq 1 - \epsilon \quad \text{and} \quad \rho_{\mu^{\text{dis}}}(\mathcal{G}_{\text{dis}}) \geq 1 - \epsilon. \quad (4.26)$$

Finally, the left inequality applies to every ergodic $\mu \in \mathfrak{G}_\beta$ when $\beta > \beta_t$, while the right inequality applies to every ergodic $\mu \in \mathfrak{G}_\beta$ when $\beta < \beta_t$.

This theorem settles, somewhat more apparently than Theorem 3 of [11], the controversy in the physics literature about whether this system does or does not undergo a first-order transition as the temperature varies; see [10] for more discussion and relevant references. The proof of Theorem 4.4 is fairly technical and it is therefore deferred to Sect. 5.

4.5 Magnetostriction transition.

Our final example is the magnetostriction transition studied recently by Shlosman and Zagreb-
nov [21]. The specific system considered in [21] has the Hamiltonian

$$H(\sigma, r) = - \sum_{\langle x, y \rangle} J(r_{x, y}) \sigma_x \sigma_y + \kappa \sum_{\langle x, y \rangle} (r_{x, y} - R)^2 + \lambda \sum_{\substack{\langle x, y \rangle, \langle z, y \rangle \\ |x - z| = \sqrt{2}}} (r_{x, y} - r_{z, y})^2. \quad (4.27)$$

Here the sites $x \in \mathbb{Z}^d$ label the atoms in a crystal; the atoms have magnetic moments represented by the Ising spins σ_x . The crystal is not rigid; the variables $r_{x, y} \in \mathbb{R}$, $r_{x, y} > 0$, play the role of spatial distance between neighboring crystal sites.

The word *magnetostriction* refers to the phenomenon where a solid undergoes a magnetic transition accompanied by a drastic change in the crystalline structure. In [21] such a transition was proven for interaction potentials $J = J(r_{x, y})$ that are strong at short distances and weak at large distances. The relevant states are characterized by disjoint *contracted*,

$$\mathcal{G}^{\text{contr}} = \left\{ (r, \sigma) : r_{x, y} \leq \eta, \forall x, y \in \Lambda_1, |x - y| = 1 \right\}, \quad (4.28)$$

and *expanded*,

$$\mathcal{G}^{\text{exp}, \pm} = \left\{ (r, \sigma) : r_{x, y} \geq \eta + \epsilon, \forall x, y \in \Lambda_1, |x - y| = 1 \right\} \cap \left\{ \sigma_x = \pm 1, \forall x \in \Lambda_1 \right\}, \quad (4.29)$$

block events. The parameters η and ϵ can be chosen so that there exists $\beta_t \in (0, \infty)$ for which the following holds:

- (1) For all $\beta \leq \beta_t$ there exists an *expanded* Gibbs state $\mu^{\text{exp}} \in \mathfrak{G}_\beta$ such that $\rho_{\mu^{\text{exp}}}(\mathcal{G}^{\text{exp}}) \geq 3/4$;
(2) For all $\beta \geq \beta_t$ there exist two distinct *contracted* Gibbs states $\mu^{\text{contr}, \pm} \in \mathfrak{G}_\beta$ such that $\rho_{\mu^{\text{contr}, \pm}}(\mathcal{G}^{\text{contr}, \pm}) \geq 3/4$.

In particular at $\beta = \beta_t$ there exist three distinct Gibbs states; one expanded and two contracted with opposite values of the magnetization. The authors conjecture that these are the only shift-ergodic Gibbs states at $\beta = \beta_t$.

Unfortunately, the above system has unbounded spins and interactions and so it is not strictly of the form for which Theorem 2.5 applies. Instead we will use the following generalization:

Theorem 4.5 *Consider a spin system with translation-invariant finite-range interaction potentials $(\Phi_A)_{A \in \mathbb{Z}^d}$ such that the torus measure is reflection positive for all even L . Let $\mathcal{G}_1, \dots, \mathcal{G}_n$ be a collection of good B -block events satisfying the requirements in Theorem 2.5 and let \mathcal{B} be the corresponding bad event. Then for all $\epsilon > 0$ there exists $\delta > 0$ such that for all $\beta \geq 0$ for which $\mathfrak{p}_\beta(\mathcal{B}) < \delta$ the following is true: If $\mu \in \mathcal{G}_\beta$ is a B -shift invariant Gibbs state with*

$$\sum_{\substack{A: A \in \mathbb{Z}^d \\ 0 \in A}} E_\mu(|\Phi_A|) < \infty, \quad (4.30)$$

then we have

$$\rho_\mu(\mathcal{B}) \in [0, \epsilon] \quad (4.31)$$

and

$$\rho_\mu(\mathcal{G}_i) \in [0, \epsilon] \cup [1 - \epsilon, 1], \quad (4.32)$$

for all $i = 1, \dots, n$.

Proof. The proof is virtually identical to that of Theorem 2.5 with one exception: Since the interactions are not bounded, we cannot use Lemma 3.4 directly. Suppose we have a Gibbs state μ that obeys (4.30) but violates one of the conditions (4.31–4.32). Let $R_{M,N}(\mathcal{C}_N)$ be as in (3.4). Lemma 3.1 still applies and so we have (3.5) for some N .

Let $L = MNB$ and let \mathcal{D}_M be the event that the boundary energy in the box Λ is less than cM^{d-1} , i.e.,

$$\mathcal{D}_M = \left\{ \sum_{\substack{A: A \cap \Lambda_L \neq \emptyset \\ A \cap \Lambda_L^c \neq \emptyset}} |\Phi_A| \leq cM^{d-1} \right\}. \quad (4.33)$$

where c is a positive constant. In light of the condition (4.30), the fact that the interaction has a finite range, and the Chebyshev bound, it is clear that we can choose c so that $\mu(\mathcal{D}_M^c) < (4N^d)^{-1}$ for all M . Hence, we have

$$\mu(\mathcal{D}_M \cap \{R_{M,N}(\mathcal{C}_N) > 1/2\}) > \frac{1}{4N^d}. \quad (4.34)$$

Next let s and s' be as in the proof of Lemma 3.4 and suppose that both s and s' belong to \mathcal{D}_M . Then, by definition,

$$|H_{\Lambda_L}(s) - H_{\Lambda_L}(s')| \leq 2cM^{d-1} \quad (4.35)$$

and, applying the rest of the proof of Lemma 3.4, we thus have

$$\mu(\mathcal{D}_M \cap \{R_{M,N}(\mathcal{C}_N) > 1/2\}) \leq e^{2\beta cM^{d-1}} \mathbb{P}_{2L,\beta}(\mathcal{D}_M \cap \{R_{M,N}(\mathcal{C}_N) > 1/2\}). \quad (4.36)$$

Neglecting \mathcal{D}_L on the right-hand side and invoking (3.28), we again derive the desired contradiction once M is sufficiently large. \square

With Theorem 4.5 in the hand, we can extract the desired conclusion for the magnetostriction transition. First, the energy condition is clearly satisfied in any state generated by tempered boundary conditions. We then know that, in every such ergodic state μ , only a small number blocks will feature bonds that are neither contracted (and magnetized) nor expanded (and non-magnetized):

$$\rho_\mu(\mathcal{G}^{\text{exp}}), \rho_\mu(\mathcal{G}^{\text{exp},\pm}) \in [0, \epsilon] \cup [1 - \epsilon, 1] \quad \text{and} \quad \rho_\mu(\mathcal{B}) \leq \epsilon. \quad (4.37)$$

The existence of a phase transition follows by noting that the contracted states have less energy than the expanded ones; there is thus a jump in the energy density as the temperature varies.

5. APPENDIX

The goal of this section is to prove Theorem 4.4 which concerns non-linear vector model with interaction (4.21). The technical part of the proof is encapsulated into the following claim:

Proposition 5.1 *There exists a constant $C_0 > 0$ such that for all $\delta > 0$ and all $C \geq C_0$ the following holds: There exists $p_0 > 0$ such that for all $p \geq p_0$ we have*

$$\sup_{\beta \geq 0} \mathfrak{p}_\beta(\mathcal{B}) < \delta \quad (5.1)$$

and

$$\lim_{\beta \rightarrow \infty} \mathfrak{p}_\beta(\mathcal{G}_{\text{dis}}) = 0 \quad \text{and} \quad \lim_{\beta \downarrow 0} \mathfrak{p}_\beta(\mathcal{G}_{\text{so}}) < \delta. \quad (5.2)$$

To prove this proposition, we will need to carry out a sequence of energy and entropy bounds. To make our energy estimates easier, and uniform in p , we first notice that there are constants $0 < a < b$ such that

$$e^{-bx^2} \leq \frac{1 + \cos(x)}{2} \leq e^{-ax^2}, \quad -1 \leq x \leq 1. \quad (5.3)$$

The argument commences by splitting \mathcal{B} into two events: The event \mathcal{B}_{wo} that Λ_1 contains a weakly-ordered bond, and $\mathcal{B}_{\text{mix}} = \mathcal{B} \setminus \mathcal{B}_{\text{wo}}$ which, as a moment's thought reveals, is the event that Λ_1 contains two adjacent bonds one of which is strongly ordered and the other disordered. The principal chessboard estimate yields the following lemma:

Lemma 5.2 *Suppose that $C \leq \sqrt{p}$. Then*

$$\mathfrak{p}_\beta(\mathcal{B}_{\text{wo}}) \leq 4 \left(\min \left\{ \frac{C^2}{\kappa} e^{-2\beta[e^{-bx^2/C^2} - e^{-a/C^2}]}, \frac{C}{\pi\sqrt{p}} e^{2\beta e^{-a/C^2}} \right\} \right)^{1/4} \quad (5.4)$$

and

$$\mathfrak{p}_\beta(\mathcal{B}_{\text{mix}}) \leq 4 \left(\min \left\{ e^{-2\beta[\frac{3}{2}e^{-b/C^2} - 1 - e^{-aC^2}]}, e^{2\beta \left(\frac{1}{\pi C\sqrt{p}}\right)^{3/4}} \right\} \right)^{1/2} \quad (5.5)$$

for all $\beta \geq 0$ and all $\kappa \in (0, 1)$. Moreover, we have

$$\mathfrak{p}_\beta(\mathcal{G}_{\text{dis}}) \leq \pi C\sqrt{p} \exp\{-2\beta[e^{-\frac{b}{C^2}} - e^{-aC^2}]\} \quad (5.6)$$

and

$$\mathfrak{p}_\beta(\mathcal{G}_{\text{so}}) \leq \frac{1}{\pi} \frac{e^{2\beta}}{C\sqrt{p}}. \quad (5.7)$$

Proof. Let Z_L be the partition function obtained by integrating $e^{-\beta H_L}$ over all allowed configurations. Consider the following reduced partition functions:

- (1) Z_L^{dis} , obtained by integrating $e^{-\beta H_L}$ subject to the restriction that every bond in \mathbb{T}_L is disordered.
- (2) Z_L^{so} , obtained similarly while stipulating that every bond in \mathbb{T}_L is strongly ordered.
- (3) Z_L^{wo} , in which every bond in \mathbb{T}_L is asked to be weakly ordered.
- (4) Z_L^{mix} , enforcing that every other horizontal line contains only strongly-ordered bonds, and the remaining lines contain only disordered bonds. A similar periodic pattern is imposed on vertical lines as well.

To prove the lemma, we will need upper and lower bounds on the partition functions in (1-2), and upper bounds on the partition functions in (3-4).

We begin by upper and lower bounds on Z_L^{dis} . First, using the fact that the Hamiltonian is always non-positive, we have $e^{-\beta H_L} \geq 1$. On the other hand, the inequalities (5.3) and a natural monotonicity of the interaction imply that

$$\left(\frac{1 + \cos(\phi_x - \phi_y)}{2} \right)^p \leq \left(\frac{1 + \cos(C/\sqrt{p})}{2} \right)^p \leq e^{-aC^2} \quad (5.8)$$

whenever $\langle x, y \rangle$ is a disordered bond. In particular, $-\beta H_L$ is less than $2\beta e^{-aC^2} |\mathbb{T}_L|$ for every configuration contributing to Z_L^{dis} . Using these observations we now easily derive that

$$(2\pi)^{|\mathbb{T}_L|} \leq Z_L^{\text{dis}} \leq (2\pi)^{|\mathbb{T}_L|} e^{2\beta e^{-aC^2} |\mathbb{T}_L|}. \quad (5.9)$$

Similarly, for the partition function Z_L^{so} get

$$\left(e^{2\beta e^{-b\kappa^2/C^2}} \frac{2\kappa}{C\sqrt{p}} \right)^{|\mathbb{T}_L|} \leq Z_L^{\text{so}} \leq 2\pi e^{2\beta |\mathbb{T}_L|} \left(\frac{2}{C\sqrt{p}} \right)^{|\mathbb{T}_L|-1}. \quad (5.10)$$

Indeed, for the upper bound we first note that $-\beta H_L \leq 2\beta |\mathbb{T}_L|$. Then we fix a tree spanning all vertices of \mathbb{T}_L , disregard the constraints everywhere except on the edges in the tree and, starting from the “leaves,” we sequentially integrate all site variables. (Thus, each site is effectively forced into an interval of length $\frac{2}{C\sqrt{p}}$, except for the “root” which retains all of its 2π possibilities.) For the lower bound we fix a number $\kappa \in (0, 1)$ and restrict the integrals to configurations such that $|\phi_x - \phi_y| \leq \frac{\kappa}{C\sqrt{p}}$ for all bonds $\langle x, y \rangle$ in \mathbb{T}_L . The bound $-\beta H_L \geq 2\beta e^{-b\kappa^2/C^2} |\mathbb{T}_L|$ then permits us to estimate away the Boltzmann factor for all configurations; the entropy factor reflects the fact that each site can vary throughout an interval of length at least $\frac{2\kappa}{C\sqrt{p}}$.

Next we will derive good upper bounds on the remaining two partition functions. First, similar estimates as those leading to the upper bound in (5.10) give us

$$Z_L^{\text{wo}} \leq 2\pi \left(e^{2\beta e^{-a/C^2}} \frac{2C}{\sqrt{p}} \right)^{|\mathbb{T}_L|}. \quad (5.11)$$

For the partition function Z_L^{mix} we note that $1/4$ of all sites are adjacent only to disordered bonds, while the remaining $3/4$ are connected to one another via a grid of strongly-ordered bonds. Estimating $-\beta H_L \leq \beta(1 + e^{-aC^2})|\mathbb{T}_L|$ for all relevant configuration, similar calculations as those leading to (5.10) again give us

$$Z_L^{\text{mix}} \leq 2\pi e^{\beta(1+e^{-aC^2})|\mathbb{T}_L|} (2\pi)^{\frac{|\mathbb{T}_L|}{4}} \left(\frac{2}{C\sqrt{p}}\right)^{\frac{3}{4}|\mathbb{T}_L|-1}. \quad (5.12)$$

It now remains to combine these estimates into the bounds on the quantities on the left-hand side of (5.4–5.5) and (5.6–5.7).

We begin with the bound (5.6). Clearly, $\mathfrak{p}_\beta(\mathcal{G}_{\text{dis}})$ is the $L \rightarrow \infty$ limit of $(Z_L^{\text{dis}}/Z_L)^{1/|\mathbb{T}_L|}$, which using the lower bound $Z_L \geq Z_L^{\text{so}}$ with $\kappa = 1$ easily implies (5.6). The bound (5.7) is obtained similarly, except that now we use that $Z_L \geq Z_L^{\text{dis}}$. The remaining two bounds will conveniently use the fact that for two-dimensional nearest-neighbor models, and square tori, the torus measure $\mathbb{P}_{L,\beta}$ is reflection positive even with respect to the diagonal planes in \mathbb{T}_L . Indeed, focusing on (5.4) for a moment, we first note that \mathcal{B}_{wo} is covered by the union of four (non-disjoint) events characterized by the position of the weakly-ordered bond on Λ_1 . If $\mathcal{B}_{\text{wo}}^{(1)}$ is the event that the lower horizontal bond is the culprit, the subadditivity property of \mathfrak{p}_β —see Lemma 6.3 of [4]—gives us $\mathfrak{p}_\beta(\mathcal{B}_{\text{wo}}) \leq 4\mathfrak{p}_\beta(\mathcal{B}_{\text{wo}}^{(1)})$. Disseminating $\mathcal{B}_{\text{wo}}^{(1)}$ using reflections in coordinate directions, we obtain an event enforcing weakly-ordered bonds on every other horizontal line. Next we apply a reflection in a diagonal line of even parity to make this into an even parity grid. From the perspective of reflections in odd-parity diagonal lines—i.e., those not passing through the vertices of the grid—half of the “cells” enforces all four bonds therein to be weakly ordered, while the other half does nothing. Applying chessboard estimates for these diagonal reflections, we get rid of the latter cells. The result of all these operations is the bound

$$\mathfrak{p}_\beta(\mathcal{B}_{\text{wo}}) \leq \lim_{L \rightarrow \infty} 4 \left(\frac{Z_L^{\text{wo}}}{Z_L}\right)^{\frac{1}{4|\mathbb{T}_L|}}. \quad (5.13)$$

Estimating Z_L from below by the left-hand sides of (5.9–5.10) now directly implies (5.4).

The event \mathcal{B}_{mix} is handled similarly: First we fix a position of the ordered-disordered pair of bonds and use subadditivity of \mathfrak{p}_β to enforce the *same* choice at every lattice plaquette; this leaves us with four overall choices. Next we use diagonal reflections to produce the event underlying Z_L^{mix} . Estimating Z_L from below by $1/4$ -th power of the lower bound in (5.9) and $3/4$ -th power of the lower bound in (5.10) with $\kappa = 1$, we get the first term in the minimum in (5.5). To get the second term, we use that $Z_L \geq Z_L^{\text{dis}}$, apply (5.12) and invoke the bound $1 + e^{-aC^2} \leq 2$. \square

Proof of Proposition 5.1. The desired properties are simple consequences of the bounds in Lemma 5.2. Indeed, if C is so large that $e^{-b/C^2} > e^{-aC^2}$, then (5.6) implies that $\mathfrak{p}_\beta(\mathcal{G}_{\text{dis}}) \rightarrow 0$ as $\beta \rightarrow \infty$. On the other hand, (5.7) shows that the $\beta \rightarrow \infty$ limit of $\mathfrak{p}_\beta(\mathcal{G}_{\text{so}})$ is order $1/\sqrt{p}$, which can be made as small as desired by choosing p sufficiently large.

To prove also (5.1), we first invoke Lemma 6.3 of [4] one last time to see that $\mathfrak{p}_\beta(\mathcal{B}) \leq \mathfrak{p}_\beta(\mathcal{B}_{\text{wo}}) + \mathfrak{p}_\beta(\mathcal{B}_{\text{mix}})$. We thus have to show that both $\mathfrak{p}_\beta(\mathcal{B}_{\text{wo}})$ and $\mathfrak{p}_\beta(\mathcal{B}_{\text{mix}})$ can be made arbitrary small by increasing p appropriately. We begin with $\mathfrak{p}_\beta(\mathcal{B}_{\text{mix}})$. Let C be so large that

$$\frac{3}{2}e^{-b/C^2} - 1 - e^{-aC^2} > 0. \quad (5.14)$$

Then for β such that $e^{2\beta} > p^{1/4}$ the first term in the minimum in (5.6) decays like a negative power of p , while for the complementary values of β , the second term is $O(p^{-1/8})$. As to the remaining term, $\mathfrak{p}_\beta(\mathcal{B}_{\text{wo}})$, here we choose $\kappa \in (0, 1)$ such that

$$e^{-b\kappa^2/C^2} - e^{-a/C^2} > 0, \quad (5.15)$$

and apply the first part of the minimum in (5.4) for β with $e^{2\beta} \geq \sqrt{p}$, and the second part for the complementary β , to show that $\mathfrak{p}_\beta(\mathcal{B}_{\text{wo}})$ is also bounded by constants time a negative power of p , independently of β . Choosing p large, (5.1) follows. \square

Now we can finally prove Theorem 4.4:

Proof of Theorem 4.4. We will plug into our main theorem. First, it is easy to check that the good block events \mathcal{G}_{so} and \mathcal{G}_{dis} satisfy the conditions (1) and (2) of Theorem 2.5. Then (5.1) and (2.15–2.16) imply (4.24–4.25). The limits (5.2) and Corollary 2.6 then imply the existence of the transition temperature β_t and of the corresponding coexisting states. Since the negative of the energy density undergoes a jump at β_t from values $\gtrsim e^{-b/C^2}$ to values $\lesssim e^{-aC^2}$ —which differ by almost one once $C \gg 1$ —all ergodic states for $\beta > \beta_t$ must have small energy density while the states for $\beta < \beta_t$ will have quite a lot of energy. Applying (4.24–4.25), all ergodic $\mu \in \mathfrak{G}_\beta$ for $\beta > \beta_t$ must be dominated by strongly-order bonds, while those for $\beta < \beta_t$ must be dominated by disordered bonds. \square

ACKNOWLEDGMENTS

The research of M.B. was supported by the NSF grant DMS-0306167 and that of R.K. by the grants GAČR 201/03/0478 and MSM 0021620845. Large parts of this paper were written while both authors visited Microsoft Research in Redmond.

REFERENCES

- [1] M. Biskup, L. Chayes, and N. Crawford, *Mean-field driven first-order phase transitions in systems with long-range interactions*, submitted.
- [2] M. Biskup, L. Chayes, and S.A. Kivelson, *Order by disorder, without order, in a two-dimensional spin system with $O(2)$ -symmetry*, Ann. Henri Poincaré **5** (2004), no. 6, 1181–1205.
- [3] M. Biskup, L. Chayes, and R. Kotecký, *Coexistence of partially disordered/ordered phases in an extended Potts model*, J. Statist. Phys. **99** (2000), no. 5/6, 1169–1206.
- [4] M. Biskup, L. Chayes, and Z. Nussinov, *Orbital ordering in transition-metal compounds: I. The 120-degree model*, Commun. Math. Phys. **255** (2005) 253–292.
- [5] M. Biskup, L. Chayes, and Z. Nussinov, *Orbital ordering in transition-metal compounds: II. The orbital-compass model*, in preparation.
- [6] L. Chayes, R. Kotecký, and S. B. Shlosman, *Aggregation and intermediate phases in dilute spin systems*, Commun. Math. Phys. **171** (1995) 203–232.
- [7] L. Chayes, R. Kotecký, and S. B. Shlosman, *Staggered phases in diluted systems with continuous spins*, Commun. Math. Phys. **189** (1997) 631–640.
- [8] L. Chayes, S. Shlosman, and V. Zagrebnov, *Discontinuity in magnetization in diluted $O(n)$ -Models*, J. Statist. Phys. **98** (2000) 537–549.
- [9] R.L. Dobrushin and S.B. Shlosman, *Phases corresponding to minima of the local energy*, Selecta Math. Soviet. **1** (1981), no. 4, 317–338.

- [10] A.C.D. van Enter and S.B. Shlosman, *First-order transitions for n -vector models in two and more dimensions: Rigorous proof*, Phys. Rev. Lett. **89** (2002) 285702.
- [11] A.C.D. van Enter and S.B. Shlosman, *Provable first-order transitions for nonlinear vector and gauge models with continuous symmetries*, Commun. Math. Phys. **255** (2005) 21–32.
- [12] J. Fröhlich, R. Israel, E. H. Lieb, and B. Simon, *Phase transitions and reflection Positivity. I. General theory and long range models*, Commun. Math. Phys. **62** (1978) 1–34.
- [13] J. Fröhlich, R. Israel, E. H. Lieb, and B. Simon, *Phase transitions and reflection positivity. II. Lattice systems with short range and Coulomb interactions*, J. Statist. Phys. **22** (1980) 297–347.
- [14] H.-O. Georgii, *Gibbs Measures and Phase Transitions*, de Gruyter Studies in Mathematics, vol. 9, Walter de Gruyter & Co., Berlin, 1988.
- [15] R. Kotecký, L. Laanait, A. Messenger and J. Ruiz, *The q -state Potts model in the standard Pirogov-Sinai theory: surface tensions and Wilson loops*, J. Statist. Phys. **58** (1990), no. 1-2, 199–248.
- [16] R. Kotecký and S.B. Shlosman, *First-order phase transitions in large entropy lattice models*, Commun. Math. Phys. **83** (1982), no. 4, 493–515.
- [17] L. Laanait, A. Messenger, S. Miracle-Solé, J. Ruiz and S. Shlosman, *Interfaces in the Potts model. I. Pirogov-Sinai theory of the Fortuin-Kasteleyn representation*, Commun. Math. Phys. **140** (1991), no. 1, 81–91.
- [18] D.H. Martirosian, *Translation invariant Gibbs states in the q -state Potts model*, Commun. Math. Phys. **105** (1986), no. 2, 281–290.
- [19] S.A Pirogov and Ya.G. Sinai, *Phase diagrams of classical lattice systems* (Russian), Theor. Math. Phys. **25** (1975) no. 3, 358–369.
- [20] S.A Pirogov and Ya.G. Sinai, *Phase diagrams of classical lattice systems. Continuation* (Russian), Theor. Math. Phys. **26** (1976), no. 1, 61–76.
- [21] S. Shlosman and V. Zagrebnov, *Magnetostriction transition*, J. Statist. Phys. **114** (2004) 563–574.
- [22] M. Zahradník, *An alternate version of Pirogov-Sinai theory*, Commun. Math. Phys. **93** (1984) 559–581.