

KINETICALLY CONSTRAINED SPIN MODELS

ABSTRACT. Kinetically constrained spin models (KSCM) are interacting particle systems with Glauber-like dynamics, no static interactions beyond hard core and reversible w.r.t. simple i.i.d product measure. The essential feature is that the creation/destruction of a particle on x can occur only if the configuration around x satisfies certain constraints (which completely define each specific model). KSCM have been introduced and studied in physical literature as examples of models showing a slow relaxation and in the attempt to model liquid/glass transition. Because of the constraints KSCM may show ergodicity breakdown (dynamical phase transition) without any equilibrium counterpart and it is this phenomenon that has been the main object of investigation in the physical community. In these lectures, starting from the basics of reversible continuous time Markov chains, we will develop rigorous tools to analyze the density and volume dependence of the relaxation time for a fairly general class of KSCM.

1. TENTATIVE PLAN

- Lecture 1:** Continuous time reversible Markov chains with finite state space. Mixing and relaxation times, spectral gap, logarithmic Sobolev constant, geometric tools and examples [10, 4, 5]
- Lecture 2:** KSCM [9, 2, 3, 13]. Definition, graphical construction of the generator and semigroup, stationary and reversible measures, ergodicity and mixing [8, 7]. Bootstrap percolation models [1, 11, 6, 12] and KSCM.
- Lecture 3:** KSCM: analysis of the relaxation time and of the persistence function [3].

REFERENCES

- [1] M. Aizenman and J. L. Lebowitz. Metastability effects in bootstrap percolation. *J. Phys. A*, 21(19):3801–3813, 1988.
- [2] David Aldous and Persi Diaconis. The asymmetric one-dimensional constrained Ising model: rigorous results. *J. Statist. Phys.*, 107(5-6):945–975, 2002.
- [3] Nicoletta Cancrini, Fabio Martinelli, Cyril Roberto, and Cristina Toninelli. Rigorous results for kinetically constrained spin models. preprint, 2006.
- [4] P. Diaconis and L. Saloff-Coste. Logarithmic Sobolev inequalities for finite Markov chains. *Ann. Appl. Probab.*, 6(3):695–750, 1996.
- [5] Persi Diaconis and Laurent Saloff-Coste. Comparison theorems for reversible Markov chains. *Ann. Appl. Probab.*, 3(3):696–730, 1993.
- [6] Alexander E. Holroyd. Sharp metastability threshold for two-dimensional bootstrap percolation. *Probab. Theory Related Fields*, 125(2):195–224, 2003.
- [7] George Kordzakhia and Steven Lalley. Ergodicity and mixing properties of the northeast models. preprint, 2006.
- [8] Thomas M. Liggett. *Interacting particle systems*. Springer-Verlag, New York, 1985.
- [9] F. Ritort and P. Sollich. Glassy dynamics of kinetically constrained models. *Advances in Physics*, 52(4):219–342, 2003.

- [10] Laurent Saloff-Coste. Lectures on finite Markov chains. In *Lectures on probability theory and statistics (Saint-Flour, 1996)*, pages 301–413. Springer, Berlin, 1997.
- [11] Roberto H. Schonmann. On the behavior of some cellular automata related to bootstrap percolation. *Ann. Probab.*, 20(1):174–193, 1992.
- [12] Cristina Toninelli and Giulio Biroli. A new class of cellular automata with a discontinuous transition. preprint, 2005.
- [13] Cristina Toninelli, Giulio Biroli, and Daniel S. Fisher. Cooperative behavior of kinetically constrained lattice gas models of glassy dynamics. *J. Stat. Phys.*, 120(1-2):167–238, 2005.